



LARVIVA ProStart CO-FEEDING DIET AND EARLY WEANING FEED

Ingredients:	Fish meal, krill meal, hyd DL-methionin, probiotic	<i>,</i> , , , , , , , , , , , , , , , , , ,	sh ge	latin, lecithin, algae, bet	taine, yeast extract
Composition:	Crude proteins	67 %		vitE added	700 mg/kg
	Crude lipids	12 %		vitA added	40,000 IU/kg
	Crude ash	11.5 %		vitD3 added	1,400 IU/kg
	Crude cellulose	0.1 %		Phosphorus	1.70 %
	vitC added	1,500 mg/kg		n-3 HUFA	2.30 %
	Depending on producing factory and composition may vary compared to a				
Sizes:	 LARVIVA ProStart 10 LARVIVA ProStart 200 LARVIVA ProStart 300 	0: 125 - 250 µ			
Sizes:	LARVIVA ProStart 20	0: 125 - 250 μ 0: 250 - 400 μ			

PRODUCT FEATURES

Agglomerated, high protein larval feed with the right amino acid balance, for co-feeding with live feed and for early weaning.

DOM. Opened bags should be used within one month and stored well closed in a refrigerator.

This sophisticated production technology allows us to produce rounded feed particles with good buoyancy in the water column. This results in a better availability to the young larvae requiring specific and complete nutrition. A special coating of the feed ensures high palatability without compromising on digestibility. Feeding can start very early as of mouth opening, co-fed with live feed.

When using LARVIVA ProStart, weaning can be done earlier and faster so the need for live feed can be significantly reduced. This results in a more cost effective and predictable larval output.

Compared to live feed, LARVIVA ProStart is available in the appropriate size range, has the complete nutritional profile and is of constant quality always off-the-shelf available.

LARVIVA ProStart includes Bactocell®, a probiotic that is documented to reduce vertebral deformities in fish larvae.

For optimal results at start feeding combine with LARVIVA Multigain enriched rotifers and/or Artemia. After weaning proceed with LARVIVA ProWean.

PRODUCT USE

Best results are obtained by feeding many times small quantities. In the beginning it is advised to feed by hand in excess and observe carefully if all larvae are feeding. Do not mix LARVIVA ProStart with water before feeding.

- LARVIVA ProStart 100: From day of first feeding, feed from 4 to 100 g/m³/day
- LARVIVA ProStart 200: 10-250 g/m³/day
- LARVIVA ProStart 300: 30-200 g/m³/day
- LARVIVA ProStart 400: 30-400 g/m³/day

Check regularly to ensure that there is enough feed in the water column. In case of overfeeding, increase water exchange to avoid pollution by uneaten feed.